arXiv:math/0701138v2 [math.AG] 23 Jun 2008

Smooth n-dimensional subvarieties of $\mathbb{P}^{2n-1}$ containing a family of very degenerate divisors

José Carlos Sierra *

June 23, 2008

Abstract

A classification and a detailed geometric description are given for smooth n-dimensional subvarieties $X \subset \mathbb{P}^{2n-1}$ containing a family of effective divisors each of them spanning a linear $\mathbb{P}^n$ of $\mathbb{P}^{2n-1}$. Some results on multisecant lines to threefolds $X \subset \mathbb{P}^5$ follow, as a byproduct.

Keywords: low codimensional subvarieties, hypersurface fibrations, Zak's Theorem on Tangencies, linear normality, multisecant lines.

MSC 2000: 14M07, 14N05, (14D06).

1 Introduction

Let $X \subset \mathbb{P}^r$ be a smooth irreducible n-dimensional projective subvariety defined over the complex numbers. If $r \geq 2n+1$, then $X \subset \mathbb{P}^r$ can be embedded in $\mathbb{P}^{2n+1}$ by a linear projection, so $X \subset \mathbb{P}^r$ is said to be a low codimensional subvariety if $r \leq 2n$. Low codimensional subvarieties are quite special. In the early 70's, Barth and Larsen showed in [4], [5] and [21] that for $i \leq 2n-r$ the cohomology groups $H^i(X, \mathbb{C})$ and the homotopy groups $\pi_i(X)$ are the same as those of $\mathbb{P}^r$ (and hence of complete intersections by Lefschetz hyperplane theorem). According to these results, Hartshorne [11] made several conjectures on low codimensional subvarieties following

---

[*] Partially supported by the MCYT project BFM2003-03971/MATE and the Santander-Complutense project PR27/05-13876.

the principle that the lower the codimension of $X \subset \mathbb{P}^r$, the closer are its properties to those of complete intersections. In particular, he conjectured that $X \subset \mathbb{P}^r$ is a complete intersection if $2r < 3n$ and that $X \subset \mathbb{P}^r$ is linearly normal if $2r-2 < 3n$. While the first question remains open, Zak [38] proved Hartshorne's conjecture on linear normality by means of the so-called Zak's Theorem on Tangencies, characterizing also in [39] the four Severi varieties as the only non-linearly normal subvarieties $X \subset \mathbb{P}^r$ in the limit case $2r-2 = 3n$.

In this paper we study low codimensional subvarieties containing a family of degenerate divisors by using Zak's Theorem on Tangencies. Barth-Larsen results imply that the Picard group of $X \subset \mathbb{P}^r$ is generated by its hyperplane section if $r \leq 2n-2$. In this range, no effective divisor D on X is contained in a linear s-dimensional subspace of $\mathbb{P}^r$ for $n-1 \leq s \leq r-2$ (see Corollary 2.5). So we start by considering the boundary case $X \subset \mathbb{P}^{2n-1}$, and the simplest situation $D = \mathbb{P}^{n-1}$.

There exist some examples of smooth subvarieties $X \subset \mathbb{P}^{2n-1}$ containing a linear divisor $D = \mathbb{P}^{n-1}$ and realizing any possible degree allowed by theory (see [6], Ex. 5.2). Moreover the only smooth subvariety $X \subset \mathbb{P}^{2n-1}$ ruled by a family of linear divisors is the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^{n-1} \subset \mathbb{P}^{2n-1}$, as Lanteri and Turrini showed [20].

This fact leads us to study possible generalizations. First, one is tempted to classify smooth subvarieties $X \subset \mathbb{P}^{2n}$ ruled by a family of linear divisors. A complete classification was given by Lanteri [19] and Aure [2] for $n = 2$, by Toma [36] for $n = 3$, and by Ionescu and Toma [16] for $n \leq 4$. The problem is still open for $n \geq 5$.

Another possibility is to classify smooth subvarieties $X \subset \mathbb{P}^{2n-1}$ containing a family of effective divisors on X each of them spanning a linear $\mathbb{P}^n$ of $\mathbb{P}^{2n-1}$. This is what we do in this note. Examples of such varieties were given by Ionescu and Toma in [15], where it is shown, for every integer $d \geq 1$, the existence of what they called (degree d)-hypersurface fibrations $X \subset \mathbb{P}^{2n-1}$ over $\mathbb{P}^1$. In this paper we prove that these are the only hypersurface fibrations in $\mathbb{P}^{2n-1}$, and we also describe the exceptional cases in which the family of hypersurfaces does not induce a fibration on X. The precise result is the following.

Theorem 3.14 Let $X \subset \mathbb{P}^{2n-1}$ be a smooth n-dimensional subvariety containing a family $\{D_t\}_{t \in T}$ of hypersurfaces of degree $d \geq 2$.

(i) if $D_t \cap D_{t'} \neq \emptyset$, then $n \leq 3$ with equality if and only if $X \subset \mathbb{P}^5$ is linked to a linear $\mathbb{P}^3$ by the complete intersection of a rank-4 quadric cone and a hypersurface of $\mathbb{P}^5$ of degree d;

(ii) if $D_t \cap D_{t'} = \emptyset$, then $Y = \bigcup_{t\in T}\langle D_t\rangle \subset \mathbb{P}^{2n-1}$ is a rational normal scroll of n-planes $S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-1}$ of degree $n-1$. Furthermore:

  (a) if $n \geq 3$, then $X \subset \mathbb{P}^{2n-1}$ is linked to a rational normal scroll of $(n-1)$-planes $S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-3}$ of degree $n-2$ by the complete intersection of $Y \subset \mathbb{P}^{2n-1}$ and a hypersurface of $\mathbb{P}^{2n-1}$ of degree $d+1$;

  (b) if $n = 2$, then $X \subset \mathbb{P}^3$ is a surface of degree $d+1$ containing a line $L \subset \mathbb{P}^3$ and $\{D_t\}_{t\in T}$ is the pencil $|H - L|$ of plane curves of degree d on X.

To obtain the proof of Theorem 3.14 we consider two cases separately. If the intersection of any two divisors of the family is non-empty, we strongly use Zak's Theorem on Tangencies to get the bound $n \leq 3$. On the other hand, if the intersection of any two divisors of the family is empty, we prove that $X \subset \mathbb{P}^{2n-1}$ is linear normal by using Zak's circle of ideas on secant varieties (see Proposition 3.8).

To complete the overview on related problems, we would like to stress that smooth n-dimensional subvarieties $X \subset \mathbb{P}^{2n}$ containing a family of hypersurfaces are still far from being classified, even for small n. As in the case of subvarieties containing a family of linear divisors, this is due to the existence of irregular subvarieties of $\mathbb{P}^{2n}$, in contrast to the case $r = 2n-1$. Consider first $n = 2$, i.e. surfaces in $\mathbb{P}^4$ containing a family of plane curves. There exists a complete classification only if the plane curves form a fibration on the surface. Ellia and Sacchiero [8] classified conic fibrations (see also [7]), and Ranestad [28] classified surfaces ruled by a family of plane curves of degree greater than or equal to three. If the plane curves do not form a fibration, just some partial results have been recently obtained in [33]. As far as we know, the case $n \geq 3$ remains unexplored. We refer to [32] for a more detailed account on low codimensional subvarieties containing a family of hypersurfaces.

The last part of the paper is devoted to obtaining some consequences on multisecant lines to a smooth threefold $X \subset \mathbb{P}^5$ by using Theorem 3.14. Extending results of Severi [31] and Aure [3] on trisecant lines to surfaces in $\mathbb{P}^4$, Kwak [18] proved that a smooth threefold $X \subset \mathbb{P}^5$ has no apparent triple points if and only if $H^0(\mathcal{I}_X(2)) \neq 0$. A classification of threefolds with no apparent quadruple points is still unknown. We present a different proof of Kwak's result (see Theorem 4.3) as well as some remarks on the structure of 4-secant lines to a threefold $X \subset \mathbb{P}^5$, strengthening both [23], Th. 2.3, and [18], Prop. 4.9. Finally, we point out that Theorem 3.14 can also be used to complete the proof of Th. 1.4 in [24] (see Remark 4.6).

2 Preliminaries

Let $X \subset \mathbb{P}^r$ be an irreducible non-degenerate (i.e. not contained in a hyperplane) projective subvariety of dimension $n \geq 2$ defined over the complex numbers.

Definition 2.1. Let D be an effective Weil divisor on $X \subset \mathbb{P}^r$, and let $\langle D \rangle = \mathbb{P}^s \subset \mathbb{P}^r$ denote its linear span. We call D a *degenerate divisor on* X if $n-1 \leq s \leq r-2$. Moreover D is said to be *linear* if $D = \mathbb{P}^{n-1}$, and D is said to be a *hypersurface* if $\langle D \rangle = \mathbb{P}^n$. Linear divisors and hypersurfaces will be called *very degenerate divisors on* X.

Let T be an irreducible projective variety. We denote by $\{D_t\}_{t \in T}$ the algebraic family of divisors on X corresponding to a closed codimension one subvariety $\mathcal{D} \subset X \times T$ flat over T. Subvarieties $X \subset \mathbb{P}^r$ containing a 2-dimensional family of very degenerate divisors are well known. They are described in Lemma 2.2 and Theorem 2.3 below.

Lemma 2.2. *Let* $X \subset \mathbb{P}^r$ *be an* n*-dimensional subvariety containing a family* $\{D_t\}_{t \in T}$ *of linear divisors on* X*. If* $\dim T \geq 2$*, then* $X = \mathbb{P}^n$*.*

Proof. Since $\dim T \geq 2$, for any two points $x, y \in X$ there exists some $t \in T$ such that $x, y \in D_t = \mathbb{P}^{n-1}$. Therefore the line $\langle x, y \rangle$ is contained in X, so $X = \mathbb{P}^n$. □

Theorem 2.3. *Let* $X \subset \mathbb{P}^r$ *be an* n*-dimensional subvariety containing a family* $\{D_t\}_{t \in T}$ *of hypersurfaces. If* $r \geq n+2$*, then* $\dim T \leq 2$*. Moreover if* $\dim T = 2$*, then* $\{D_t\}_{t \in T}$ *is a family of (maybe reducible) quadric hypersurfaces and* $X \subset \mathbb{P}^r$ *is one of the following subvarieties:*

(i) the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$*;*

(ii) the rational normal scroll $S_{1,2} \subset \mathbb{P}^4$*;*

(iii) the Veronese surface $v_2(\mathbb{P}^2) \subset \mathbb{P}^5$ or its projection to $\mathbb{P}^4$;

(iv) a cone with vertex $\mathbb{P}^{n-2}$ over a curve;

(v) a cone over varieties (i), (ii), (iii).

Proof. If $n = 2$, the result was given by C. Segre in [30]. If $n \geq 3$, we reduce to the case $n = 2$ by intersecting $X \subset \mathbb{P}^r$ with $n - 2$ general hyperplanes. If this intersection is a cone with vertex a point over a curve, then it is easy to see that $X \subset \mathbb{P}^r$ is a cone with vertex $\mathbb{P}^{n-2}$ over a curve. We conclude by recalling that the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$ is the only non-trivial extension of $S_{1,2} \subset \mathbb{P}^4$, and that neither the Veronese surface $v_2(\mathbb{P}^2) \subset \mathbb{P}^5$ nor its projection to $\mathbb{P}^4$ can be non-trivially extended. These facts follow, for instance, from [37] and [34]. $\square$

From now on we also assume that $X \subset \mathbb{P}^r$ is smooth (except in Section 2.1). The following result is a corollary of Barth-Larsen theorems [4], [5] and [21].

Theorem 2.4 (Barth-Larsen)**.** *Let $X \subset \mathbb{P}^r$ be a smooth subvariety of dimension n.*

(i) if $r \leq 2n - 1$, then $H^1(X, \mathcal{O}_X) = 0$;

(ii) if $r \leq 2n - 2$, then the restriction $\mathrm{Pic}(\mathbb{P}^r) \to \mathrm{Pic}(X)$ determines an isomorphism.

Corollary 2.5. *Let $X \subset \mathbb{P}^r$ be a smooth subvariety of dimension n. Let D be a degenerate divisor on X. Then $r \geq 2n - 1$.*

Proof. Let H denote the hyperplane divisor on $X \subset \mathbb{P}^r$. Since $D \subset X$ is a degenerate divisor, there exists an effective non-zero divisor $E \in |H - D|$ on X. If $r \leq 2n-2$, then $\mathrm{Pic}(X) \cong \mathbb{Z}$ generated by H. Therefore $D \equiv_{\mathrm{lin}} aH$ and $E \equiv_{\mathrm{lin}} bH$ for some positive integers a, b. Hence $H \equiv_{\mathrm{lin}} D + E \equiv_{\mathrm{lin}} (a + b)H$, yielding a contradiction (cf. Remark 2.11). $\square$

The bound $r \geq 2n - 1$ is sharp, as the following example shows.

Example 2.6 (Beltrametti-Howard-Schneider-Sommese)**.** We present an example of smooth n-dimensional subvarieties $X \subset \mathbb{P}^{2n-1}$ containing a linear $\mathbb{P}^{n-1}$ (see [6], Ex. 5.2). Fix integers $s \geq 0$ and $n \geq 2$. Let $\mathcal{P} =$

$\mathbb{P}(\mathcal{O}_{\mathbb{P}^{2n-1}}(1) \oplus \mathcal{O}_{\mathbb{P}^{2n-1}}(s+1))$ and let $p : \mathcal{P} \to \mathbb{P}^{2n-1}$ denote the bundle projection. Let ξ denote the tautological line bundle on $\mathcal{P}$. The intersection of n general elements of $|\xi|$ is a smooth n-dimensional subvariety $X' \subset \mathcal{P}$ which is mapped isomorphically by p onto its image $X \subset \mathbb{P}^{2n-1}$. The embedding $\mathcal{P} \subset \mathbb{P}^{\dim|\xi|}$ given by ξ contains a linear subspace $\mathbb{P}^{2n-1}$ corresponding to $\mathbb{P}(\mathcal{O}_{\mathbb{P}^{2n-1}}(1))$, so $X' \subset \mathcal{P}$ (and hence $X \subset \mathbb{P}^{2n-1}$) contains a linear $\mathbb{P}^{n-1}$. It also follows that $N_{\mathbb{P}^{n-1}/X} \cong \mathcal{O}_{\mathbb{P}^{n-1}}(-s)$. If $s = 0$, then $\deg X = n$ and $X \subset \mathbb{P}^{2n-1}$ is the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^{n-1} \subset \mathbb{P}^{2n-1}$. If $s \geq 1$, then $\deg X = \frac{(s+1)^n - 1}{s}$. These are the only possible degrees of smooth n-dimensional subvarieties $X \subset \mathbb{P}^{2n-1}$ containing a linear $\mathbb{P}^{n-1}$, as it is showed in [6], Prop. 8.1.

We focus on smooth subvarieties $X \subset \mathbb{P}^{2n-1}$ containing a family $\{D_t\}_{t\in T}$ of linear divisors.

Remark 2.7. Let $X \subset \mathbb{P}^r$ be a smooth n-dimensional subvariety containing a family $\{D_t\}_{t\in T}$ of linear divisors on X. If $D_t \cap D_{t'} \neq \emptyset$ for any $t, t' \in T$, Lemma 3.3 below yields $D_t \cap D_{t'} = \mathbb{P}^{n-2}$. Then it is straightforward to see that $X = \mathbb{P}^n$ (otherwise $X \subset \mathbb{P}^r$ is a cone of vertex $\mathbb{P}^{n-2}$, whence singular). Therefore, we can assume $D_t \cap D_{t'} = \emptyset$.

Smooth subvarieties $X \subset \mathbb{P}^{2n-1}$ ruled by a family $\{D_t\}_{t\in T}$ of linear divisors were first characterized in [20].

Theorem 2.8 (Lanteri-Turrini)**.** *The only smooth n-dimensional subvariety of $\mathbb{P}^{2n-1}$ ruled by a family of linear divisors is the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^{n-1} \subset \mathbb{P}^{2n-1}$.*

In Section 3, we pass on to classify smooth n-dimensional subvarieties $X \subset \mathbb{P}^{2n-1}$ containing a family $\{D_t\}_{t\in T}$ of hypersurfaces, i.e. effective divisors on X such that $\langle D_t \rangle = \mathbb{P}^n$ for every $t \in T$. First, we include a few words regarding the main tool we use to obtain our classification.

2.1 Zak's Theorem on Tangencies

The geometry of projective subvarieties is well reflected in the relative position of their embedded tangent spaces. Modern algebraic geometry reduced basically the study of tangents spaces to some computations involving Chern classes. Fortunately, a new dimension in the understanding of projective varieties by means of the relative position of the tangent spaces was introduced by F.L. Zak and his celebrated theorem on tangencies.

Definition 2.9. Let $T_xX \subset \mathbb{P}^r$ denote the embedded tangent space to $X \subset \mathbb{P}^r$ at $x \in X$. A linear subspace $L \subset \mathbb{P}^r$ is said to be *tangent to* $X \subset \mathbb{P}^r$ *along* $Z \subset X$, if $T_xX \subset L$ for every $x \in Z$. We denote $T(Z, X) := \bigcup_{x \in Z} T_xX \subset \mathbb{P}^r$.

Theorem 2.10 (Zak's Theorem on Tangencies)**.** *Let* $X \subset \mathbb{P}^r$ *be a non-degenerate subvariety of dimension* n*. If a linear subspace* $L \subsetneq \mathbb{P}^r$ *is tangent to* $X \subset \mathbb{P}^r$ *along a closed subvariety* $Z \subset X$*, then*

$$\dim Z \leq \dim L - n.$$

Theorem 2.10 leads to the proof of Hartshorne's conjecture on linear normality (see [40] II, Cor. 2.17). Finiteness of Gauss mappings and the bound on the dimension of the dual variety ([40] I, Cor. 2.8 and Cor. 2.5) also follow as an immediate consequence. References for interesting applications of Theorem 2.10 by other authors can be found in [40]. See [22] for a brief account on this topic.

Remark 2.11. Corollary 2.5, which is the starting point of the paper, can be obtained from Theorem 2.10 as well (see [40] I, Prop. 2.16). This second proof does not depend on Theorem 2.4.

3 Subvarieties $X \subset \mathbb{P}^{2n-1}$ containing a family of hypersurfaces

Let $X \subset \mathbb{P}^{2n-1}$ be a smooth n-dimensional subvariety containing a family $\{D_t\}_{t \in T}$ of hypersurfaces. Denote $\mathbb{P}^n_t := \langle D_t \rangle \subset \mathbb{P}^{2n-1}$ and $Y := \bigcup_{t \in T} \mathbb{P}^n_t \subset \mathbb{P}^{2n-1}$.

Remark 3.1. If $X \subset \mathbb{P}^{2n-1}$ is not the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$, then Theorem 2.3 yields $\dim T = 1$ and $\dim Y = n + 1$.

Lemma 3.2. *Let* $X \subset \mathbb{P}^{2n-1}$ *be a smooth subvariety of dimension* n*. Let* $\{D_t\}_{t \in T}$ *be a family of effective divisors on* X*. Then* D_t *and* $D_{t'}$ *are linearly equivalent divisors on* X *for any* $t, t' \in T$*.*

Proof. It follows from Theorem 2.4 that $H^1(X, \mathcal{O}_X) = 0$. Then D_t is algebraically equivalent to $D_{t'}$ if and only if D_t is linearly equivalent to $D_{t'}$. $\square$

3.1 Case $D_t \cap D_{t'} \neq \emptyset$

The next statement, that has already been quoted in Remark 2.7, can be found, for instance, in [10], Th. 17.24.

Lemma 3.3. *Let Z_1, Z_2 be pure dimensional subvarieties of a smooth variety X. Then the codimension of their intersection (if non-empty) is at most the sum of their codimensions.*

Lemma 3.3 yields $n-2 \leq \dim(\mathbb{P}^n_t \cap \mathbb{P}^n_{t'}) \leq n-1$ for general $t, t' \in T$. We analyze both cases separately.

Lemma 3.4. *Let $X \subset \mathbb{P}^{2n-1}$ be a smooth subvariety of dimension n. Let $\{D_t\}_{t\in T}$ be a family of hypersurfaces such that $D_t \cap D_{t'} \neq \emptyset$. If*

$$\dim(\mathbb{P}^n_t \cap \mathbb{P}^n_{t'}) = n-1$$

for general $t, t' \in T$, then $n = 2$.

Proof. If $\dim(\mathbb{P}^n_t \cap \mathbb{P}^n_{t'}) = n-1$ for general $t, t' \in T$, then either $Y = \mathbb{P}^{n+1}$ (and hence $n = 2$) or $Y \subset \mathbb{P}^{2n-1}$ is a cone with vertex $\mathbb{P}^{n-1}$. In the latter case, it follows from Remark 3.1 that $\dim Y = n+1$. Let $y \in Y$ be a general point and let $T_yY = \mathbb{P}^{n+1} \subset \mathbb{P}^{2n-1}$ be the tangent space to Y at y. Then $T_yY \subset \mathbb{P}^{2n-1}$ is tangent to Y along $\langle y, \mathbb{P}^{n-1}\rangle \backslash \mathbb{P}^{n-1}$. Therefore $T_yY \subset \mathbb{P}^{2n-1}$ is tangent to X along $X_y := \langle y, \mathbb{P}^{n-1}\rangle \cap X$. Since $\dim X_y = n-1$, it follows from Theorem 2.10 that $n \leq 2$. ☐

Lemma 3.5. *Let $X \subset \mathbb{P}^{2n-1}$ be a smooth subvariety of dimension n. Let $\{D_t\}_{t\in T}$ be a family of hypersurfaces such that $D_t \cap D_{t'} \neq \emptyset$. If*

$$\dim(\mathbb{P}^n_t \cap \mathbb{P}^n_{t'}) = n-2$$

for general $t, t' \in T$, then $n = 3$.

Proof. Fix general $t, t' \in T$. Since $\dim(\mathbb{P}^n_t \cap \mathbb{P}^n_{t'}) = n-2$, then $D_t \cap D_{t'} = \mathbb{P}^{n-2}$. If $X \subset \mathbb{P}^{2n-1}$ is not the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$, the family $\{D_t\}_{t\in T}$ of hypersurfaces is the pencil of divisors on X generated by D_t and $D_{t'}$. This follows from Remark 3.1 and Lemma 3.2. In particular, $\mathbb{P}^{n-2} \subset D_t$ for every $t \in T$ and $Y \subset \mathbb{P}^{2n-1}$ is an $(n+1)$-dimensional cone with vertex $\mathbb{P}^{n-2}$. As before, the tangent space $T_yY = \mathbb{P}^{n+1} \subset \mathbb{P}^{2n-1}$ to Y at a general $y \in Y$ is tangent to X along $X_y := \langle y, \mathbb{P}^{n-2}\rangle \cap X$. Since $\dim X_y = n-2$, Theorem 2.10 yields $n \leq 3$. Therefore $n = 3$. ☐

Notation 3.6. Let $a_0 \leq \ldots \leq a_n$ be a non-trivial sequence of non-negative integers. We denote by $S_{a_0,\ldots,a_n} \subset \mathbb{P}^{n+a_0+\cdots+a_n}$ the rational normal scroll of n-planes of degree $a_0 + \cdots + a_n$. For $E = \oplus_{i=0}^n \mathcal{O}_{\mathbb{P}^1}(a_i)$, we can also see $S_{a_0,\ldots,a_n} \subset \mathbb{P}^{n+a_0+\cdots+a_n}$ as the image of $\mathbb{P}(E)$ in $\mathbb{P}^{n+a_0+\cdots+a_n}$ by the tautological line bundle $\mathcal{O}_{\mathbb{P}(E)}(1)$ on $\mathbb{P}(E)$. Recall that the rational normal scroll of n-planes $S_{1,\ldots,1} \subset \mathbb{P}^{2n+1}$ of degree $n+1$ is precisely the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^n \subset \mathbb{P}^{2n+1}$.

Proposition 3.7. *Let $X \subset \mathbb{P}^{2n-1}$ be a smooth n-dimensional subvariety containing a family $\{D_t\}_{t\in T}$ of hypersurfaces such that $D_t \cap D_{t'} \neq \emptyset$. Then $n \leq 3$, with equality if and only if $X \subset \mathbb{P}^5$ is linked to a linear $\mathbb{P}^3$ by the complete intersection of a rank-4 quadric cone and a hypersurface of $\mathbb{P}^5$.*

Proof. Lemmas 3.4 and 3.5 yield $n \leq 3$. The Segre embedding $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$ is linked to a linear $\mathbb{P}^3$ by the complete intersection of a rank-4 quadric cone and a quadric hypersurface of $\mathbb{P}^5$. If $X \subset \mathbb{P}^5$ is not this Segre embedding, it follows from Remark 3.1 and the proof of Lemma 3.5 that $Y \subset \mathbb{P}^5$ is a rational scroll of 3-planes of vertex $L = \mathbb{P}^1$. Since $X \subset \mathbb{P}^5$ is linearly normal by [38], and $D_t \cap D_{t'} = L$, it follows that $Y \subset \mathbb{P}^5$ is a rational normal scroll of 3-planes. Hence $\deg Y = 2$ and $Y \subset \mathbb{P}^5$ is the quadric cone $S_{0,0,1,1} \subset \mathbb{P}^5$. The result follows, for instance, from [17], Th. 2.1. □

3.2 Case $D_t \cap D_{t'} = \emptyset$

In this case, the main ingredient of Theorem 3.14 is the following proposition.

Proposition 3.8. *Let $X \subset \mathbb{P}^{2n-1}$ be a smooth subvariety of dimension n. Let $\{D_t\}_{t\in T}$ be a family of hypersurfaces. If $D_t \cap D_{t'} = \emptyset$, then $X \subset \mathbb{P}^{2n-1}$ is linearly normal.*

Proof. Suppose to the contrary that $X \subset \mathbb{P}^r$ is a non-degenerate n-dimensional subvariety that can be isomorphically projected to $\mathbb{P}^{2n-1}$. We denote by $SX \subset \mathbb{P}^r$ the secant variety of $X \subset \mathbb{P}^r$. Let $z \in SX$ be a general point and let $E_z \subset X$ be the corresponding entry locus, i.e. the set of points in X swept out by secant lines to $X \subset \mathbb{P}^r$ passing through z. Since $\dim SX = 2n-1$, then $\dim E_z = 2$. Terracini's Lemma [35] yields $T_xX \subset T_zSX$ for every $x \in E_z$. Let $J(D_t, D_{t'}) \subset \mathbb{P}^r$ denote the join of D_t and $D_{t'}$, i.e. the set of points contained in lines spanned by a point of D_t and a point of $D_{t'}$. We claim that $J(D_t, D_{t'})$ has the expected dimension

$$\dim J(D_t, D_{t'}) = 2(n-1) + 1 = 2n-1.$$

Otherwise, projecting to $\mathbb{P}^{2n-2}$ from a suitable linear subspace of $\mathbb{P}^r$, we get $\emptyset = D_t \cap D_{t'} \subset \mathbb{P}^{2n-2}$, which is not possible. For every $t \in T$, denote $Z_t = E_z \cap D_t$. Since $\dim J(D_t, D_{t'}) = \dim SX$, it follows that $J(D_t, D_{t'}) = SX$ for any $t, t' \in T$, $t \neq t'$. Therefore, for every $t \in T$, there exist infinitely many secant lines to X passing through z and meeting D_t. Thus $\dim Z_t \geq 1$ for every $t \in T$, and so $\dim Z_t = 1$ for a general $t \in T$, since $\dim E_z = 2$. Theorem 2.10 yields $T(Z_t, D_t) = \mathbb{P}^n_t \subset \mathbb{P}^{2n-1}$, and hence

$$Y = \bigcup_{t\in T} \mathbb{P}^n_t = \bigcup_{t\in T} T(Z_t, D_t) \subset \bigcup_{t\in T} T(Z_t, X) = T(E_z, X) \subset T_z SX,$$

so $Y \subset T_z SX$. But $T_z SX \subset \mathbb{P}^r$ is a linear subspace of dimension $2n-1$. Hence $X \subset \mathbb{P}^r$ is a degenerate subvariety, a contradiction. □

Lemma 3.9. *Let $X \subset \mathbb{P}^{2n-1}$ be a smooth subvariety of dimension n. Let $\{D_t\}_{t\in T}$ be a family of hypersurfaces. If $D_t \cap D_{t'} = \emptyset$, then $Y \subset \mathbb{P}^{2n-1}$ is a rational normal scroll of n-planes $S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-1}$ of degree $n-1$.*

Proof. $X \subset \mathbb{P}^{2n-1}$ is a linearly normal subvariety by Proposition 3.8. It follows from Theorem 2.3 and Lemma 3.2 that T is a rational curve. Since $D_t \cap D_{t'} = \emptyset$, we deduce that $Y \subset \mathbb{P}^{2n-1}$ is a rational normal scroll of n-planes. Therefore $\deg Y = n-1$. We claim that $\dim(\mathbb{P}^n_t \cap \mathbb{P}^n_{t'}) = 1$ for any $t, t' \in T$. Otherwise, if $\dim(\mathbb{P}^n_t \cap \mathbb{P}^n_{t'}) \geq 2$, then $D_t \cap D_{t'} \neq \emptyset$. Thus we obtain $Y = S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-1}$. □

Example 3.10 (Ionescu-Toma)**.** For every integer $d \geq 1$, the existence of smooth n-dimensional (degree d)-hypersurface fibrations $X \subset \mathbb{P}^{2n-1}$ contained in $Y = S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-1}$ was shown in [15], Prop. 3. We reproduce their construction (some notations have been changed). Let $E = \mathcal{O}_{\mathbb{P}^1}^{\oplus 2} \oplus \mathcal{O}_{\mathbb{P}^1}(1)^{\oplus n-1}$ and let $\mathbb{P}(E)$ be the projective bundle over $\mathbb{P}^1$. Let $\pi : \mathbb{P}(E) \to \mathbb{P}^1$ be the projection, $H = \mathcal{O}_{\mathbb{P}(E)}(1)$ and $F = \pi^*\mathcal{O}_{\mathbb{P}^1}(1)$. It follows from Bertini's theorem that the general element $X \in |dH + F|$ is smooth. Let $\varphi : \mathbb{P}(E) \to Y \subset \mathbb{P}^{2n-1}$ be the morphism induced by H. Then the restriction $\varphi : X \to \mathbb{P}^{2n-1}$ turns out to be an embedding onto its image. We prove in the sequel that moreover elements $X \in |dH + F|$ are the only smooth divisors on $\mathbb{P}(E)$ such that $\varphi : X \to \mathbb{P}^{2n-1}$ is an embedding.

Proposition 3.11. *With the above notations, let $X \in |dH + bF|$ be a smooth divisor on $\mathbb{P}(E)$. If $\varphi : X \to \mathbb{P}^{2n-1}$ is an embedding, $X \notin |F|$, then $b = 1$.*

Proof. Set $E' = \mathcal{O}_{\mathbb{P}^1}^{\oplus 2}$, $E'' = \mathcal{O}_{\mathbb{P}^1}(1)^{\oplus n-1}$ and $S = \mathbb{P}(E')$. Let $p_1 = \pi_{|S}$ and $p_2 = \varphi_{|S}$ be the two projections of $S \cong \mathbb{P}^1 \times \mathbb{P}^1$. Since S is mapped by φ onto the vertex $L = \mathbb{P}^1$ of Y, it follows that $\varphi : \mathbb{P}(E) \backslash S \to Y \backslash L \subset \mathbb{P}^{2n-1}$ is an embedding. Therefore $\varphi : X \to \mathbb{P}^{2n-1}$ is an embedding only if the restriction $\varphi : X \cap S \to L$ is an embedding. Then $C = X \cap S$ is a rational curve on S and $C \in |dF_1 + bF_2|$, where $F_i = p_i^*(\mathcal{O}_{\mathbb{P}^1}(1))$. Since φ contracts F_1, we get $1 = F_1 \cdot C = b$. □

Remark 3.12. To get a more projective flavor, we present a different geometric description of $X \in |dH + F|$ embedded in $Y \subset \mathbb{P}^{2n-1}$. Let $R \in |H - F|$. Since $H \equiv_{\text{lin}} R + F$, it follows that $dH + F \equiv_{\text{lin}} (d+1)H - R$. If $n \geq 3$, then $\varphi(R) \subset Y$ is a rational normal scroll of $(n-1)$-planes $S_{0,0,1,\ldots,1} \subset \mathbb{P}^{2n-3}$ of degree $n-2$. If $n = 2$, then $\varphi(R) = L \subset \mathbb{P}^3$. Rational normal scrolls are projectively normal (see [1], Prop. 2.9), i.e. the restriction maps

$$H^0(\mathcal{O}_{\mathbb{P}^{2n-1}}(m)) \to H^0(\mathcal{O}_Y(m))$$

are surjective for every integer $m \geq 1$. Therefore if $n \geq 3$, then $X \subset \mathbb{P}^{2n-1}$ is linked to a rational normal scroll of $(n-1)$-planes $S_{0,0,1,\ldots,1} \subset \mathbb{P}^{2n-3}$ of degree $n-2$ by the complete intersection of Y and a hypersurface of $\mathbb{P}^{2n-1}$ of degree $d+1$. On the other hand, if $n = 2$, then $X \subset \mathbb{P}^3$ is a surface of degree $d+1$ containing $L = \mathbb{P}^1$, and $\{D_t\}_{t \in T}$ is the pencil $|H - L|$ of plane curves of degree d on X.

Remark 3.13. Let us look more closely at $n = 3$. In this case $Y = S_{0,0,1,1} \subset \mathbb{P}^5$ is a rank-4 quadric cone. Hence $Y \subset \mathbb{P}^5$ is ruled by two pencils of linear $\mathbb{P}^3$'s, say T_1 and T_2. If, for instance, $X \subset \mathbb{P}^5$ is linked to a $\mathbb{P}^3$ of T_2 by the complete intersection of Y and a hypersurface of $\mathbb{P}^5$ of degree $d+1$, then $\{D_t\}_{t \in T_1}$ induces a fibration on X by hypersurfaces of degree d. On the other hand, $\{D_t\}_{t \in T_2}$ is a pencil of hypersurfaces of degree $d+1$ with base locus the vertex $L = \mathbb{P}^1$ of $Y \subset \mathbb{P}^5$, as in Proposition 3.7.

The main result of the paper is now obtained.

Theorem 3.14. *Let $X \subset \mathbb{P}^{2n-1}$ be a smooth n-dimensional subvariety containing a family $\{D_t\}_{t \in T}$ of hypersurfaces of degree $d \geq 2$.*

(i) if $D_t \cap D_{t'} \neq \emptyset$, then $n \leq 3$ with equality if and only if $X \subset \mathbb{P}^5$ is linked to a linear $\mathbb{P}^3$ by the complete intersection of a rank-4 quadric cone and a hypersurface of $\mathbb{P}^5$ of degree d;

(ii) if $D_t \cap D_{t'} = \emptyset$, then $Y = \bigcup_{t\in T}\langle D_t\rangle \subset \mathbb{P}^{2n-1}$ is a rational normal scroll of n-planes $S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-1}$ of degree $n-1$. Furthermore:

  (a) if $n \geq 3$, then $X \subset \mathbb{P}^{2n-1}$ is linked to a rational normal scroll of $(n-1)$-planes $S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-3}$ of degree $n-2$ by the complete intersection of $Y \subset \mathbb{P}^{2n-1}$ and a hypersurface of $\mathbb{P}^{2n-1}$ of degree $d+1$;

  (b) if $n = 2$, then $X \subset \mathbb{P}^3$ is a surface of degree $d+1$ containing a line $L \subset \mathbb{P}^3$ and $\{D_t\}_{t\in T}$ is the pencil $|H-L|$ of plane curves of degree d on X.

Proof. If follows from Propositions 3.7, 3.8, 3.11, Lemma 3.9 and Remark 3.12. □

For $n \geq 3$, the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^{n-1} \subset \mathbb{P}^{2n-1}$ is linked to a rational normal scroll of $(n-1)$-planes $S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-3}$ of degree $n-2$ by the complete intersection of $Y \subset \mathbb{P}^{2n-1}$ and a quadric hypersurface of $\mathbb{P}^{2n-1}$. Thus we can state Theorems 2.8 and 3.14 together if $n \geq 3$.

Corollary 3.15. *Let $X \subset \mathbb{P}^{2n-1}$ be a smooth n-dimensional subvariety containing a family of very degenerate divisors on X. If $n \geq 3$, then $X \subset \mathbb{P}^{2n-1}$ is linked to a rational normal scroll of $(n-1)$-planes $S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-3}$ of degree $n-2$ by the complete intersection of a rational normal scroll of n-planes $S_{0,0,1,\dots,1} \subset \mathbb{P}^{2n-1}$ of degree $n-1$ and a hypersurface of $\mathbb{P}^{2n-1}$.*

Remark 3.16. Proposition 3.11 can be obtained as a particular case of [12], Th. 3.7. In fact, using the terminology of [12], we can reformulate Corollary 3.15 as follows: a smooth n-dimensional subvariety $X \subset \mathbb{P}^{2n-1}$ contains a family of very degenerate divisors if and only if $X \subset \mathbb{P}^{2n-1}$ is a linearly normal *Roth variety.* We wish to thank Paltin Ionescu for pointing out this paper.

Remark 3.17. In particular, smooth n-dimensional subvarieties $X \subset \mathbb{P}^{2n-1}$ containing a family of hypersurfaces are linearly normal. This is no longer true if either $X \subset \mathbb{P}^{2n}$ or $\langle D_t\rangle = \mathbb{P}^{n+1}$, as the following example shows.

Example 3.18. The Veronese surface $v_2(\mathbb{P}^2) \subset \mathbb{P}^5$ contains a family $\{D_t\}_{t\in T}$ of conics and can be isomorphically projected to $\mathbb{P}^4$. The Segre embedding $\mathbb{P}^2 \times \mathbb{P}^2 \subset \mathbb{P}^8$ contains a family $\{D_t\}_{t\in T}$ of divisors $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$ and can be isomorphically projected to $\mathbb{P}^7$. In both cases $D_t \cap D_{t'} \neq \emptyset$. On the other hand, if $D_t \cap D_{t'} = \emptyset$, it is easy to see that subvarieties $X \subset \mathbb{P}^{2n}$ ruled by a family of linear divisors are linearly normal. We do not know if hypersurface fibrations $X \subset \mathbb{P}^{2n}$ are linearly normal as well. At least, this is known to be true for $n \leq 3$ (cf. [9]).

4 Multisecant lines to threefolds in $\mathbb{P}^5$

Let $\Sigma_k(X) \subset \mathbb{G}(1, r)$ be the variety of k-secant lines to a smooth irreducible non-degenerate projective subvariety $X \subset \mathbb{P}^r$, and let $S_k(X) \subset \mathbb{P}^r$ be the closure of the union of all k-secant lines. If $X \subset \mathbb{P}^5$ is a threefold, then $S_2(X) = \mathbb{P}^5$. Moreover, it follows from the (dimension+2)-secant lemma [27] that $S_5(X) \neq \mathbb{P}^5$. We say that $X \subset \mathbb{P}^5$ has no apparent triple (resp. quadruple) points if $S_3(X) \neq \mathbb{P}^5$ (resp $S_4(X) \neq \mathbb{P}^5$). Threefolds in $\mathbb{P}^5$ with no apparent triple points were classified by Kwak [18], extending to $n = 3$ the following result of Aure [3].

Theorem 4.1 (Aure). *Let $X \subset \mathbb{P}^4$ be a smooth surface. Then $S_3(X) \neq \mathbb{P}^4$ if and only if either $H^0(\mathcal{I}_X(2)) \neq 0$ or $X \subset \mathbb{P}^4$ is a quintic elliptic scroll.*

We reobtain Kwak's result in Theorem 4.3, as a byproduct of Theorem 3.14 and the following lemma (see [29]).

Lemma 4.2 (B. Segre). *Let $V \subset \mathbb{P}^r$ be an irreducible subvariety of dimension n. Let $\Sigma_\infty \subset \mathbb{G}(1, r)$ be a component of maximal dimension of the variety of lines contained in V. Then:*

- *(i) if $\dim \Sigma_\infty = 2n - 2$, then $V = \mathbb{P}^n$;*
- *(ii) if $\dim \Sigma_\infty = 2n - 3$, then $V \subset \mathbb{P}^r$ is either a quadric hypersurface or a scroll of linear $\mathbb{P}^{n-1}$'s over a curve.*

Theorem 4.3 (Kwak). *Let $X \subset \mathbb{P}^5$ be a smooth threefold. Then $S_3(X) \neq \mathbb{P}^5$ if and only if $H^0(\mathcal{I}_X(2)) \neq 0$.*

Proof. If $X \subset \mathbb{P}^5$ is contained in a quadric hypersurface $Q \subset \mathbb{P}^5$, then $S_3(X) \subset Q$. For the converse, let $C \subset \mathbb{P}^3$ be the intersection of $X \subset \mathbb{P}^5$ with a general $\mathbb{P}^3$ of $\mathbb{P}^5$. If $\dim \Sigma_3(X) \leq 4$, then $C \subset \mathbb{P}^3$ is a smooth curve with at most a finite number of trisecant lines. So $C \subset \mathbb{P}^3$ is either a rational normal cubic or the complete intersection of two quadrics of $\mathbb{P}^3$. Therefore

$X \subset \mathbb{P}^5$ is either the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$ or the complete intersection of two quadric hypersurfaces of $\mathbb{P}^5$. In both cases $H^0(\mathcal{I}_X(2)) \neq 0$. So we assume $\dim \Sigma_3(X) = 5$. Then $S_3(X) \subset \mathbb{P}^5$ is either a quadric hypersurface or a scroll of $\mathbb{P}^3$'s over a curve by Lemma 4.2. If the second occurs, then $S_3(X) \subset \mathbb{P}^5$ is a quadric cone $S_{0,0,1,1} \subset \mathbb{P}^5$ by Theorem 3.14. So $X \subset \mathbb{P}^5$ is contained in the quadric hypersurface $S_3(X) \subset \mathbb{P}^5$. □

Corollary 4.4. *Let $X \subset \mathbb{P}^{n+2}$ be a smooth subvariety of dimension n. Then $S_3(X) \neq \mathbb{P}^{n+2}$ if and only if either $H^0(\mathcal{I}_X(2)) \neq 0$ or $X \subset \mathbb{P}^4$ is a quintic elliptic scroll.*

Proof. Let $C \subset \mathbb{P}^3$ denote the intersection of $X \subset \mathbb{P}^{n+2}$ with $n-1$ general hyperplanes. If $\dim \Sigma_3(X) \leq 2n-2$, then $C \subset \mathbb{P}^3$ is a smooth curve with at most a finite number of trisecant lines and $X \subset \mathbb{P}^{n+2}$ is either a complete intersection of two quadric hypersurfaces or $X \subset \mathbb{P}^5$ is the Segre embedding $\mathbb{P}^1 \times \mathbb{P}^2 \subset \mathbb{P}^5$. If $\dim \Sigma_3(X) = 2n-1$, then $S_3(X) \subset \mathbb{P}^{n+2}$ is either a quadric hypersurface or a scroll of $\mathbb{P}^n$'s over a curve by Lemma 4.2. In the scroll case, $X \subset \mathbb{P}^{n+2}$ contains a family of degenerate divisors, so Corollary 2.5 yields $n \leq 3$. Thus the result follows from Theorems 4.1 and 4.3. □

A classification of threefolds $X \subset \mathbb{P}^5$ with no apparent quadruple points is still an open problem. If $H^0(\mathcal{I}_X(3)) \neq 0$, then $X \subset \mathbb{P}^5$ has no apparent quadruple points but it is not known if there exists such a threefold satisfying $H^0(\mathcal{I}_X(3)) = 0$ (see [18] and [23]). Theorem 3.14 is useful to sharpen [23], Th. 2.3.

Proposition 4.5. *Let $X \subset \mathbb{P}^5$ be a smooth threefold. If $\dim \Sigma_k(X) = 5$ for some integer $k \geq 4$, then $H^0(\mathcal{I}_X(2)) \neq 0$.*

Proof. We claim that $S_k(X) \neq \mathbb{P}^5$. To the contrary, assume $S_k(X) = \mathbb{P}^5$. Let $H \subset \mathbb{P}^4$ be a general hyperplane. Denote $S = X \cap H \subset \mathbb{P}^4$. Let $p \in H$ be a general point. Since $\dim \Sigma_k(X) = 5$, there exists a 1-dimensional family of k-secant lines to $X \subset \mathbb{P}^5$ passing through p. Thus $p \in H$ is contained in some k-secant line to $S \subset \mathbb{P}^4$, contradicting [27]. Therefore $S_k(X) \subset \mathbb{P}^5$ is either a quadric hypersurface or a scroll of $\mathbb{P}^3$'s over a curve. Hence the result follows from Theorem 3.14. □

Remark 4.6. It was stated in [24], Th. 1.4, that a smooth threefold $X \subset \mathbb{P}^5$ containing a family $\{D_t\}_{t \in T}$ of surfaces of $\mathbb{P}^3$ is contained in a quadric hypersurface of $\mathbb{P}^5$. In the proof, the case $D_t \cap D_{t'} \neq \emptyset$ was not considered.

This fact was communicated by Emilia Mezzetti to Angelo Felice Lopez, who pointed it out to the author. Thus Theorem 3.14 also helps to fill this gap.

Finally, we slightly improve Prop. 4.9 in [18] by means of Theorem 3.14.

Proposition 4.7. *Let* $X \subset \mathbb{P}^5$ *be a smooth threefold. If* $S_4(X) \neq \mathbb{P}^5$ *and* $H^0(\mathcal{I}_X(3)) = 0$*, then* $\dim \Sigma_4(X) = 4$.

Proof. If $\dim \Sigma_4(X) = 5$, then $S_4(X) \subset \mathbb{P}^5$ is either a quadric hypersurface or a scroll of $\mathbb{P}^3$'s over a curve by Lemma 4.2. Hence $S_4(X) \subset \mathbb{P}^5$ is a quadric hypersurface by Theorem 3.14, so $H^0(\mathcal{I}_X(3)) \neq 0$. Now assume $\dim \Sigma_4(X) \leq 3$. Then $\deg X \leq 8$ by the proof of Prop. 4.9 in [18]. Smooth threefolds $X \subset \mathbb{P}^5$ up to degree 8 are described in [13], [14], [25] and [26]. They are contained in a cubic hypersurface of $\mathbb{P}^5$ except Palatini scroll, which has one apparent quadruple point. Therefore $\dim \Sigma_4(X) = 4$. ☐

Remark 4.8. It was stated in [18], Prop. 4.9, that $S_4(X) \subset \mathbb{P}^5$ is a hypersurface if $H^0(\mathcal{I}_X(3)) = 0$ and $\deg X \geq 9$, but no proof is given for this assertion when $\dim \Sigma_4(X) = 4$. If $\dim \Sigma_4(X) = 4$ and $\dim S_4(X) \leq 3$, then each of the 4-dimensional irreducible components of $\Sigma_4(X) \subset \mathbb{G}(1, 5)$ sweeps out a linear $\mathbb{P}^3$ of $\mathbb{P}^5$ by Lemma 4.2. This might happen if $X \subset \mathbb{P}^5$ contains a finite number of surfaces of degree at least four, each of them contained in some $\mathbb{P}^3$.

References

[1] Arbarello, E., Harris, J.: Canonical curves and quadrics of rank 4. Compositio Math. **43** (1981), no. 2, 145–179.

[2] Aure, A.: On surfaces in projective 4–space. Ph.D. Thesis. Oslo, 1987.

[3] Aure, A.: The smooth surfaces in $\mathbb{P}^4$ without apparent triple points. Duke Math. J. **57** (1988), no. 2, 423–430.

[4] Barth, W.: Transplanting cohomology classes in complex-projective space. Amer. J. Math. **92** (1970), 951–967.

[5] Barth, W., Larsen, M.E.: On the homotopy groups of complex projective algebraic manifolds. Math. Scand. **30** (1972), 88–94.

[6] Beltrametti, M.C., Howard, A., Schneider, M., Sommese, A.J.: Projections from subvarieties. Complex analysis and algebraic geometry, 71–107, de Gruyter, Berlin, 2000.

[7] Braun, R., Ranestad, K.: Conic bundles in projective fourspace. Algebraic geometry (Catania, 1993/Barcelona, 1994), 331–339, Lecture Notes in Pure and Appl. Math., 200, Dekker, New York, 1998.

[8] Ellia, Ph., Sacchiero, G.: Smooth surfaces of $\mathbb{P}^4$ ruled in conics. Algebraic geometry (Catania, 1993/Barcelona, 1994), 49–62, Lecture Notes in Pure and Appl. Math., 200, Dekker, New York, 1998.

[9] Fujita, T.: Projective threefolds with small secant varieties. Sci. Papers College Gen. Ed. Univ. Tokyo **32** (1982), no. 1, 33–46.

[10] Harris, J.: Algebraic geometry. A first course. Corrected reprint of the 1992 original. Graduate Texts in Mathematics, 133. Springer-Verlag, New York, 1995.

[11] Hartshorne, R.: Varieties of small codimension in projective space. Bull. Amer. Math. Soc. **80** (1974), 1017–1032.

[12] Ilic, B.: Geometric properties of the double-point divisor. Trans. Amer. Math. Soc. **350** (1998), no. 4, 1643–1661.

[13] Ionescu, P.: Embedded projective varieties of small invariants. Algebraic geometry, Bucharest 1982 (Bucharest, 1982), 142–186, Lecture Notes in Math., 1056, Springer, Berlin, 1984.

[14] Ionescu, P.: Embedded projective varieties of small invariants. II. Rev. Roumaine Math. Pures Appl. **31** (1986), no. 6, 539–544.

[15] Ionescu, P., Toma, M.: Boundedness for some special families of embedded manifolds. Classification of algebraic varieties (L'Aquila, 1992), 215–225, Contemp. Math., 162, Amer. Math. Soc., Providence, RI, 1994.

[16] Ionescu, P., Toma, M.: On very ample vector bundles on curves. Internat. J. Math. **8** (1997), no. 5, 633–643.

[17] Kwak, S.: Castelnuovo-Mumford regularity bound for smooth threefolds in $\mathbb{P}^5$ and extremal examples. J. Reine Angew. Math. **509** (1999), 21–34.

[18] Kwak, S.: Smooth threefolds in $\mathbb{P}^5$ without apparent triple or quadruple points and a quadruple-point formula. Math. Ann. **320** (2001), no. 4, 649–664.

[19] Lanteri A.: On the existence of scrolls in $\mathbb{P}^4$. Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8) **69** (1980), no. 5, 223–227 (1981).

[20] Lanteri, A., Turrini, C.: Some formulas concerning nonsingular algebraic varieties embedded in some ambient variety. Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. **116** (1982), no. 3-6, 463–474 (1984).

[21] Larsen, M.E.: On the topology of complex projective manifolds. Invent. Math. **19** (1973), 251–260.

[22] Lazarsfeld, R.: Positivity in algebraic geometry. I. Classical setting: line bundles and linear series. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics, 48. Springer-Verlag, Berlin, 2004.

[23] Mezzetti, E.: On quadrisecant lines of threefolds in $\mathbb{P}^5$. Dedicated to Silvio Greco on the occasion of his 60th birthday (Catania, 2001). Matematiche (Catania) **55** (2000), no. 2, 469–481 (2002).

[24] Mezzetti, E., Portelli, D.: Threefolds in $\mathbb{P}^5$ with a 3-dimensional family of plane curves. Manuscripta Math. **90** (1996), no. 3, 365–381.

[25] Okonek, Ch.: 3-Mannigfaltigkeiten im $\mathbb{P}^5$ und ihre zugehörigen stabilen Garben. Manuscripta Math. **38** (1982), no. 2, 175–199.

[26] Okonek, Ch.: Über 2-codimensionale Untermannigfaltigkeiten vom Grad 7 in $\mathbb{P}^4$ und $\mathbb{P}^5$. Math. Z. **187** (1984), no. 2, 209–219.

[27] Ran, Z.: The (dimension+2)-secant lemma. Invent. Math. **106** (1991), no. 1, 65–71.

[28] Ranestad, K.: On smooth plane curve fibrations in $\mathbb{P}^4$. Geometry of complex projective varieties (Cetraro, 1990), 243–255, Sem. Conf., 9, Mediterranean, Rende, 1993.

[29] Segre, B.: Sulle V_n contenenti più di $\infty^{n-k}S_k$. I and II. Atti Accad. Naz. Lincei. Rend. Cl. Sci. Fis. Mat. Nat. (8) **5** (1948), 193–197 and 275–280.

[30] Segre, C.: Le superficie degli iperspazi con una doppia infinità di curve piane o spaziali. Atti Accad. Sci. Torino Cl. Sci. Fis. Mat. Natur. **56** (1920-1921), 75–89.

[31] Severi, F.: Intorno ai punti doppi impropri di una superficie generale dello spazio a quattro dimensioni e ai suoi punti tripli apparenti. Rend. Circ. Mat. Palermo **15** (1901), 33-51.

[32] Sierra, J.C.: Low codimensional subvarieties of projective spaces containing a family of very degenerate divisors, to appear in *Vector Bundles and Low Codimensional Subvarieties: State of the Art and Recent Developments* (Trento, 2006), eds. G. Casnati, F. Catanese and R. Notari, Quaderni di Matematica (Department of Mathematics, Seconda Università di Napoli, Caserta).

[33] Sierra, J.C., Tironi, A.L.: Some remarks on surfaces in $\mathbb{P}^4$ containing a family of plane curves. J. Pure Appl. Algebra **209** (2007), no. 2, 361–369.

[34] Swinnerton-Dyer, H.P.F.: An enumeration of all varieties of degree 4. Amer. J. Math. **95** (1973), no. 2, 403–418.

[35] Terracini, A.: Sulle V_k per cui la varietà degli S_h (h+1)-seganti ha dimensione minore dell' ordinario. Rend. Circ. Mat. Palermo **31** (1911), 392–396.

[36] Toma, M.: Three-dimensional scrolls in $\mathbb{P}^6$. Arch. Math. (Basel) **65** (1995), no. 5, 444–448.

[37] X.X.X.: Correspondence. Amer. J. Math. **79** (1957), no. 4, 951–952.

[38] Zak, F.L.: Projections of algebraic varieties. (Russian) Mat. Sb. (N.S.) **116(158)** (1981), no. 4, 593–602, 608.

[39] Zak, F.L.: Severi varieties. (Russian) Mat. Sb. (N.S.) **126(168)** (1985), no. 1, 115–132, 144.

[40] Zak, F.L.: Tangents and secants of algebraic varieties. Translated from the Russian manuscript by the author. Translations of Mathematical Monographs, 127. American Mathematical Society, Providence, RI, 1993.

José Carlos Sierra
Departamento de Álgebra
Facultad de Ciencias Matemáticas
Universidad Complutense de Madrid
28040 Madrid, Spain
e-mail: jcsierra@mat.ucm.es